

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Protect-us
DOMINATE™
INSECTICIDE

ACTIVE CONSTITUENT: 100 g/L FIPRONIL

GROUP 2B INSECTICIDE



For the control of ants, earwigs, millipedes and house crickets around domestic and commercial structures as specified in the Directions for Use Table.

PROTECT-US AUSTRALIA PTY LTD

ACN 156 662 216
Unit 3, 2 Duck Street
Auburn NSW 2144

www.protect-us.com.au

CUSTOMER SERVICE 1800 420 144
EMERGENCY RESPONSE (ALL HOURS) 1800 420 144

™ Trademark of Protect-us.
APVMA Approval No. 69421/112828

DIRECTIONS FOR USE

RESTRAINT

DO NOT apply to excessively wet soils, immediately after or during heavy rain, to avoid runoff of the chemical.

DO NOT apply at less than label rates.

DO NOT apply to internal surfaces.

DO NOT treat trees bearing edible fruit or nuts.

SITUATION	PEST	RATE	CRITICAL COMMENTS
External areas & surrounds of a Domestic, Commercial, Public, and Industrial buildings & structures	Nuisance ants Including but not limited to: coastal brown ants (<i>Pheidole megacephala</i>), black house ants (<i>Ochetellus glaber</i> <i>Mayr</i>), meat ants (<i>Iridomyrmex purpureus</i>), white-footed house ants (<i>Technomyrmex albipes</i>)	60 mL in 10 L of water	Apply to trails, surfaces where ants are tracking and nests. Reapply as necessary. For perimeter or harbourage treatments, apply the prepared emulsion to a band of soil or vegetation up to two to three metres wide around and adjacent to the structure. Also treat the foundation of the structure to a height of approximately 300 mm. Use a spray volume of 5 L per 100 m ² . Higher volumes of water may be needed if organic matter is present or foliage is dense. Do not exceed the point of run-off. Pay particular attention to dark protected areas such as weep holes, cracks & crevices.

SITUATION	PEST	RATE	CRITICAL COMMENTS
External areas & surrounds of a Domestic, Commercial, Public, and Industrial buildings & structures	Millipedes (<i>Ommatoiulus moreletii</i>)	60 mL in 10 L of water	Treat external surfaces 300 mm up and 300 mm out from where the building or structure touches the ground. Apply at the rate of 1 L of prepared suspension per 25 linear metres. Pay particular attention to potential entry points, such as weep holes, cracks and crevices. Also apply to where pests may be active away from the building including retaining walls, fences, garden beds, sheds etc. Apply at the rate of 1 L of prepared suspension per 16 m ² . May be applied directly to pests as a spot treatment.
	House crickets (<i>Acheta domestica</i>) Earwigs (<i>Forficula auricularia</i>)		Apply directly onto house crickets and earwigs as a spot treatment
Spot application to ant nests in domestic situations	Nesting ants, including but not limited to: meat ants (<i>Iridomyrmex purpureus</i>)		Treat the nest entrance or mound, and where ants are active away from the nest. Apply at the rate of 1 L of prepared suspension per 16 m ² or 60 mL per m ² .

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Half fill the spray tank with water and then add the required quantity of Protect-us DOMINATE Insecticide. Stir and then top up the spray tank to the required volume. The use of this product in a tank mix with other insecticides is not recommended as the behaviour and efficacy of the product may be affected. Ensure equipment is free of leaks and clean from residues of other chemicals before mixing Protect-us DOMINATE Insecticide.

Insecticide Resistance Warning

GROUP 2B INSECTICIDE

For insecticide resistance management Protect-us DOMINATE Insecticide is a Group 2B insecticide. Some naturally-occurring insect biotypes resistant to Protect-us DOMINATE Insecticide and other Group 2B insecticides may exist through normal genetic variability in any insect population. The resistant

individuals can eventually dominate the insect population if Protect-us DOMINATE Insecticide or other Group 2B insecticides are used repeatedly. The effectiveness of Protect-us DOMINATE Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Protect-us Australia Pty Ltd accepts no liability for any losses that may result from the failure of Protect-us DOMINATE Insecticide to control resistant insects.

Protect-us DOMINATE Insecticide may be subject to specific resistance management strategies. For further information contact your Protect-us Australia Pty Ltd representative or local agricultural department agronomist.

PRECAUTIONS

Residents and pets should not be allowed in a room being treated. Any spills should be cleaned up before leaving the room (refer to the SDS). Avoid application around edible plants.

RE-ENTRY PERIOD

DO NOT re-enter treated areas until spray has dried.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Highly toxic to fish and aquatic organisms. Do NOT apply to areas where surface water is present. Rinse waters and runoff from treated areas MUST be prevented from entering drains or waterways. Do NOT apply if heavy rains are expected to occur within 48 hours of application. Do NOT contaminate streams, rivers or waterways with the chemical or used containers.

Dangerous to bees.

PROTECTION OF PETS AND LIVESTOCK

Before spraying remove animals and pets from the areas to be treated. Cover or remove any open food and water containers. Cover or remove (as applicable) fish ponds, aquariums etc. before spraying.

NOTICE

Protect-us Australia Pty Ltd (Protect-us) warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with Directions for Use under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions or under off-label permits not endorsed by Protect-us, or under abnormal conditions.

